

# THE CRITICAL GROWTH RATE FOR PARTICLE INCORPORATION DURING THE DIRECTIONAL SOLIDIFICATION OF SOLAR SILICON UNDER MICROGRAVITY

**Tina Sorgenfrei<sup>1</sup>, Thomas Jauß<sup>1</sup>, Arne Cröll<sup>1</sup>, Maral Azizi<sup>2</sup>, Christian Reimann<sup>2</sup>,  
Jochen Friedrich<sup>2</sup>, Martin Volz<sup>3</sup>**

<sup>1</sup>Crystallography, University of Freiburg, Germany

<sup>2</sup> Fraunhofer IISB, Erlangen, Germany

<sup>3</sup>NASA Marshall Space Flight Center, Huntsville, AL, USA

UNI  
FREIBURG

Crystallography  
Geo- and Environmental Natural Sciences  
Albert-Ludwigs University Freiburg


# Outline

---

- Introduction
- Experiment - 1g
  - Results
- Experiment -  $\mu\text{g}$ 
  - Results
- Outlook



# Introduction



# Introduction

---

Most important particle species in solar silicon:

$\text{SiC}$ ,  $\text{Si}_3\text{N}_4$

Sources:

- Feedstock, graphite elements of the furnace, crucible material, crucible coating

Particles formed by precipitation when elements in the melt reach oversaturation



# Introduction

---


Aim of the project:

Determine the critical growth rate in directional solidified solar silicon, to control the incorporation of foreign phase particles

- Reduction of unwanted particles in the bulk crystal volume to minimize shunts and wafer loss during wafering
- Defined incorporation of getter particles to improve the electrical properties of the wafer


# Introduction

## Influencing forces on the particles:




### Forces on the particle:

- Drag of the liquid melt
- Lift force due to convectional melt movement
- Interface force
- Gravity



# Introduction

## Particle behaviour in silicon ingots for solar cells:



Experiments and images by Fraunhofer IISB

### 4 $\mu\text{m}$ &20 $\mu\text{m}$ SiC particles:

- Float to the top of the ingot
- Can agglomerate to large clusters
- Can stick to the bottom of the crucible

### 20 $\mu\text{m}$ SiC particles:

- Show precipitates in the centre of the ingot

Small particles are pushed by the solid/liquid interface

### 150 $\mu\text{m}$ SiC particles:

- Stay at the bottom of the ingot

No pushing, engulfment!



# Introduction

---

Why does the model deviate from the experiments?

Model was simplified:

- Interface roughness neglected
- Particles were assumed to be spherical
- Melt convection was not considered (no gravity!)
- Gravitational segregation (no gravity!)
- No chemical interactions in the melt assumed

Experimental conditions close to the existing  
models needed





# Experiment

---

The experiment has to provide:


- Smooth solid/liquid interface
- Defined growth rates
- Diffusive or controlled convective conditions
- Defined/known particle size, shape, and chemistry
- NO GRAVITY




# Experiment

## TEM 02-3 ELLI module

- Mono ellipsoidal mirror furnace for floating zone growth of silicon
- Option for 7mT rotating magnetic field (50Hz)
- Oxide layer on sample surface suppresses Marangoni convection (diffusive conditions, shown on TEXUS 12)




# Experiment – 1g conditions





# Results – 1g conditions

Floating zone growth of Si in the ELLI furnace:




# Results – 1g

Si [100] rod with 4mg of SiC particles (7&60 $\mu$ m size)




Using infrared and reflected light reveals:

- Size and location of particles
- Distinguishable clusters and individual particles
- Precipitates
- Twins  $\rightarrow$  can serve as locator for particles
- Striations  $\rightarrow$  show actual growth rate and phase boundaries




# Results – 1g






# Experiment – $\mu\text{g}$ conditions





# Results – $\mu\text{g}$ conditions







# Results $\mu\text{g}$ conditions



# Results $\mu g$ conditions



Depot pushed for appr. 1.2mm  
Capture velocity 2.24mm/min  
Particle size ???  $\rightarrow$  IR

Now: Input for improved models  
(J.J.Derby) and phase field  
simulations (H. Emmerich)

# Outlook

---

- Determination of particle sizes (defined data points for incorporation)
- Location of single particles (defined data points for incorporation)
- Input of capture velocities into models and phase field simulations
- Finish characterization of TEXUS 51 crystals
  
- Prepare for TEXUS 53
  - Different particle sizes/shapes/chemistry
  - Different crystal orientation (?)

